



GLYPHOSATE: UNSAFE ON ANY PLATE


FOOD TESTING RESULTS AND
SCIENTIFIC REASONS FOR CONCERN



REPORT BY **FOOD DEMOCRACY NOW!** AND **THE DETOX PROJECT**

GLYPHOSATE: UNSAFE ON ANY PLATE

ALARMING LEVELS OF MONSANTO'S GLYPHOSATE FOUND IN POPULAR AMERICAN FOODS




“For the first time in the history of the world, every human being is now subjected to contact with dangerous chemicals from the moment of conception until death...These chemicals are now stored in the bodies of the vast majority of human beings, regardless of age. They occur in the mother's milk, and probably in the tissues of the unborn child.”¹

—RACHEL CARSON, SILENT SPRING

“Glyphosate was significantly higher in humans [fed] conventional [food] compared with predominantly organic [fed] humans. Also the glyphosate residues in urine were grouped according to the human health status. Chronically ill humans had significantly higher glyphosate residues in urine than healthy humans”²

—MONIKA KRUGER, ENVIRONMENTAL & ANALYTICAL TOXICOLOGY



“Analysis of individual tissues demonstrated that bone contained the highest concentration of [¹⁴C] glyphosate equivalents (0.3–31ppm). The remaining tissues contained glyphosate equivalents at a concentration of between 0.0003 and 11ppm. In the bone and some highly perfused tissues, levels were statistically higher in males than in females.”³

—PESTICIDE RESIDUES IN FOOD, JOINT FAO/WHO MEETING 2004

- 1 Rachel Carson, Silent Spring. (Houghton Mifflin, 1961), Elixirs of Death, 15-16.
- 2 Krüger M, Schledorn P, Schrödl W, Hoppe HW, Lutz W, et al. (2014) Detection of Glyphosate Residues in Animals and Humans. J Environ Anal Toxicol 4: 210
- 3 Residues in Food, 2004, Evaluations Part II, Toxicological, Joint FAO/WHO Meeting on Pesticide Residues. http://apps.who.int/iris/bitstream/10665/43624/1/9241665203_eng.pdf

Contents

What Is in This Report?

Findings: The first ever independent, FDA-registered laboratory food testing results for glyphosate residues in iconic American food brands finds alarming levels of glyphosate contamination and reveal the inadequacy of current food safety regulations relating to allowable pesticide residues.

What's at Stake and What the Science Actually Says About Glyphosate Safety

Part I

Executive Summary	1
A Brief History of Glyphosate	2
Why Are We Being Exposed to Increasing Amounts of Glyphosate?.....	3
• Glyphosate Exposure Has Skyrocketed since Roundup Ready GMO Crops Were Adopted.....	3
Glyphosate Food Testing Results: (in parts per billion – ppb)	5
The Laboratory	7
Method for Food Testing	7
Where Else Has Glyphosate Been Found?	7
• Glyphosate Residues Found in Food, Urine, Breast Milk, Rainwater, Rivers, Tap Water and Tampons – But FDA Has Never Done Proper Testing.....	7
Pure Science: Glyphosate Damage by the Numbers (in parts per billion).....	8
What Do These Numbers Mean?.....	9

Part II

Reasons for Concern: Glyphosate Risks to Human Health	10
Where Do Claims of “Safe Levels” Come From?.....	12
Reasons for Concern: Allowed Levels of Glyphosate Are Unlikely to be Safe	12
What Have U.S. Food Safety Regulators Done to Protect Us?.....	13
• Obama Administration Raised Glyphosate Residue Levels at Monsanto’s Request	13
• Acceptable Daily Intake (ADI) or How Much Pesticide Residue You May Eat in a Day	13
• Changes in Daily Exposure Based on Industry Science, at Monsanto’s Request and a History of Scientific Fraud	14
• U.S. Acceptable Daily Intake for Glyphosate Originally Set by EPA at 0.1ppm	14
• Current Scientific Research Calls for Much Lower Allowable Glyphosate Residues for Human Food Products	15
• Monsanto and EPA Claim Roundup and Glyphosate are Perfectly Safe: Science Says Otherwise	17
• Roundup Formula 125 Times More Toxic than Glyphosate Alone	17
Glyphosate Bio-Accumulates in Major Organs and Bones	17
Peer-Reviewed Science on Glyphosate	18
How To Avoid Glyphosate	21
Call to Action	22
• Demand Immediate Release of Industry Science Data; End of Pre-Harvest Spraying on Wheat, Oats, Barley and other Food Crops	22
• Urgent Need for Fundamental Reforms of Scientific Review Process	22
Media Contacts	22



Executive Summary

A leading FDA-registered food safety testing laboratory has found extremely high levels of the pesticide glyphosate in some of America's most popular food products.

Glyphosate, the active ingredient in Monsanto's Roundup, is the most heavily used chemical weedkiller in food and agricultural production in human history, as a result of the widespread adoption of genetically engineered crops now grown on more than 175 million acres in the United States (U.S.) and more than 440 million acres around the globe.¹

New scientific evidence shows that probable harm to human health could begin at ultra-low levels of glyphosate e.g. 0.1 parts per billion (ppb). Popular foods tested for glyphosate measured between 289.47 ppb and at levels as high as 1,125.3 ppb.

The testing and analysis was performed by Anresco Laboratories, San Francisco, an FDA registered laboratory that has performed expert food safety testing since 1943. The laboratory found that well-known products tested for glyphosate, Original Cheerios, for example, measured levels as high as 1,125.3 ppb. Other high levels of glyphosate were found in familiar products such as Oreos, Doritos, and Ritz Crackers, among 29 foods tested.

Currently, U.S. regulators allow a very high level of daily glyphosate residue in America's food. The acceptable daily intake (ADI) limit is set at 1.75 milligrams per kilogram of bodyweight per day (written 1.75 mg/kg bw/day) in the U.S., versus a more cautious 0.3 mg/kg bw/day in the European Union. Tolerances have been set through the submission of corporate-sponsored studies and industry influence on the regulatory process.

New research shows that Roundup causes liver and kidney damage in rats as reflected in changes in the functions of 4,000 genes at only 0.05 parts per billion (ppb) glyphosate equivalent indicating damage.² Additional studies have found that levels as low as 10 ppb can have toxic effects on the livers of fish³ and cause significant damage to the livers and

kidneys of rats at 700 ppb,⁴ which is the allowable level of glyphosate found in U.S. drinking water.⁵

Credible independent, peer-reviewed scientific evidence now shows that the levels of harm to human health could begin at the ultra-low levels of 0.1 parts per billion (ppb) of glyphosate.

These groundbreaking new findings that one of the most iconic cereals in U.S. contains levels as high as 1,125.3 ppb should be a wake-up call for all Americans regarding unacceptable levels of pesticide residues in our nation's food. These findings are especially troubling, considering that the latest independent scientific evidence, during which a team of international scientists re-evaluated the same data previously used by regulators, calls for a much lower ADI to be set at 0.025 mg/kg of bodyweight per day or "12 times lower than the ADI"⁶ currently set in Europe and 70 times lower than the level currently allowed by the EPA in the United States.

It's important for individuals and parents to understand that glyphosate contamination cannot be removed by washing and is not broken down by cooking or baking. Glyphosate residues can remain stable in food for a year or more, even if the foods are frozen or processed.

The testing and analysis was performed at the request of [FOOD DEMOCRACY NOW!](#), in coordination with [THE DETOX PROJECT](#), which gathered additional scientific evidence from around the world and included a compendium of independent research on glyphosate that contains Anresco Laboratory's findings.

Based on this new information, [FOOD DEMOCRACY NOW!](#) is calling for a federal investigation into the likely harmful effects of glyphosate on human health and the environment and is also seeking an investigation into the relationships between the regulators and the regulated industries, which has resulted in the public being exposed to levels of glyphosate which scientific studies show can be damaging to human health.



A Brief History of Glyphosate

Glyphosate was Originally Patented to Clean Pipes, Like Drano – 1964

Glyphosate is the presumed active ingredient of Monsanto's Roundup weedkiller and other commercial glyphosate-based herbicide formulations. However, it was first patented in 1964 by Stauffer Chemical Company in Westport, Connecticut as a chelator⁷, for removing unwanted mineral deposits from metal pipes like Drano.

Monsanto Discovers Weed-killing Properties – 1974

A few years later, glyphosate was also found to be an effective herbicide by Monsanto's John E. Franz⁸ and brought to market by the St. Louis-based company in 1974 as a non-selective, water-soluble herbicide with a specific mechanism of action: the directed interruption of plant development through metabolic poisoning.

Today, generic glyphosate formulations are produced by at least 100 manufacturers and can be found in more than 750 products worldwide, with Monsanto still dominating the market with more than \$4.75 billion in sales in 2015 alone.⁹

Glyphosate, Nutrients and Disease

As with the original pipe cleaning patent, glyphosate also binds (chelates) vital nutrients such as iron, manganese, zinc, and boron in the soil, preventing plants from taking them up.^{10 11 12 13} This could have serious implications for humans, farm animals and pets that consume genetically engineered Roundup Ready crops, as it could negatively affect the nutritional value of food.

GMO soy plants treated with glyphosate have lower levels of essential nutrients and reduced growth, compared with GMO and non-GMO soy controls not treated with glyphosate.¹⁴ Lower nutrient uptake may partly account for the increased susceptibility of GMO soy to disease,¹⁵ as well as its lower yield. Additionally, glyphosate use has been linked to higher levels of soil borne pathogens, like the Fusarium fungus and sudden death syndrome in Roundup Ready GMO soybeans.¹⁶

A German-Egyptian team of researchers found that all cows tested from Danish dairy farms excreted glyphosate in their urine. Unexpectedly low levels of manganese and cobalt were observed in all animals, which the authors said could be explained due to the strong metal chelating effect of glyphosate. Potential signs of liver and kidney toxicity were also found in the cows, which the authors noted were consistent with the findings of rodent feeding studies with GMO glyphosate-tolerant plants.¹⁷

Monsanto Receives Antimicrobial Patent – 2010

In addition to mineral chelation and herbicidal modes of action or treatment, in 2003, Monsanto applied for a patent regarding the potential antibiotic or antimicrobial activity of glyphosate. The patent, finally awarded in 2010,¹⁸ has led to renewed concern from scientists that low levels of glyphosate could have an antimicrobial effect and negative impact on human health. Emerging scientific evidence suggests that glyphosate can alter human and animal intestinal flora and may lead to a harmful imbalance in the stomach's microbiome, thus causing a decrease in beneficial gut bacteria and a rise in more toxic or harmful bacteria like it does in the soil.

Why Are We Being Exposed to Increasing Amounts of Glyphosate?

Glyphosate Exposure Has Skyrocketed since Roundup Ready GMO Crops Were Adopted

Glyphosate (N-phosphonomethylglycine) is the most heavily used chemical weedkiller in food and agricultural production in human history. Since glyphosate-based herbicides were first introduced in 1974, glyphosate use has skyrocketed more than 300-fold, leading to more than 3.5 billion pounds being sold in the United States during this time. In the past decade alone, more than 2.4 billion pounds have been applied to farmers' fields - or 67% of all glyphosate ever sold in the U.S - as a result of the widespread adoption of genetically engineered crops.¹⁹

As the main active ingredient in Monsanto's bestselling herbicide Roundup, each year more than 300 million pounds of glyphosate-based herbicides are sprayed on food crops, lawns and home gardens across the country.²⁰

In the past 20 years, there's been a massive increase in glyphosate use directly linked to the introduction of Monsanto's Roundup Ready genetically engineered crops²¹, such as GMO corn, soybeans, cotton, sugar beets and canola plants that have been specifically manipulated to survive being sprayed with glyphosate-based herbicides.²²

According to U.S. Department of Agriculture (USDA) data, 93% of all soybeans and 89% corn and cotton planted by farmers in the U.S. are genetically engineered to be Roundup Ready or glyphosate tolerant.²³

In one form or another, whether as refined sugars such a high fructose corn syrup or beet sugar,


oils, vitamins or binding agents, ingredients from these genetically engineered plants are included in 75 to 80% of the processed foods sold in grocery stores across the country. Along with the increased exposure risks due to Roundup's use as a pre-harvest drying agent in wheat, oats, barley and other crops, Americans now face a continual exposure to glyphosate residues in our most popular foods.

Today the levels of Roundup and other glyphosate-based herbicides used by American farmers are so high that the U.S. Geological Survey (USGS) has found glyphosate in more than 75 percent of rain and rainwater samples across the Midwest.²⁴ This means that when farmers spray Roundup on their crops, the chemicals not only run off into local rivers and streams, but glyphosate also evaporates into the air, into local cloud cover, where it can then rain on communities downwind even hundreds of miles from the original source of application.


Overall, researchers at the USGS detected glyphosate in 60 to 100% of both the air and rain samples collected in Iowa, Mississippi and Indiana. When glyphosate is sprayed on farmers' fields it moves widely into the environment and further downstream from where the chemical was intended for agricultural use.²⁵

The USGS estimates glyphosate accounted for 53.5% of total agricultural herbicide use in 2009.²⁶ Each year, due to the increasing growth of weeds resistant to glyphosate-based herbicides, farmers are forced to spray more toxic weedkillers on their fields to combat newly evolving superweeds. Today, more than 80 million acres of U.S. farmland are covered with glyphosate-resistant superweeds. Now, with the approval new GMO crops designed to be resistant to additional herbicides like 2,4-D and dicamba, consumers will be exposed to more toxic combinations of persistent pesticide residues in their food.²⁷

Estimated Agricultural Use for Glyphosate, 1992
EPest-Low



Estimated Agricultural Use for Glyphosate, 2014 (Preliminary) EPest-High



In addition to finding high levels of glyphosate in tested food products, a small number of the foods tested by Anresco Laboratories were also contaminated with aminomethylphosphonic acid (AMPA), a metabolite of glyphosate, which is considered a weak organic acid with a phosphonic acid group and is one of the primary degradation products of glyphosate. Regulators claim that glyphosate and AMPA have similar toxicological profiles, but more research is needed to confirm AMPA's potential toxic impacts.

With an increasingly steady diet of processed foods, the American public is being primed for massive and previously unanticipated pesticide residue levels in our food, which will only continue to go up with the continued adoption and USDA approval of new herbicide-tolerant GMO crops.

In addition to on-farm use, Roundup and

glyphosate-based herbicides are also sprayed widely in parks, public spaces, lawns, home gardens, on roadsides, in forestry management, and used as a pre-harvest drying agent on certain food crops, including oats, wheat, barley and edible dry beans.²⁸

With the widespread increase in glyphosate use over the past 20 years and the fact that independent science has confirmed low level exposure to Roundup causes liver and kidney damage at only 0.05 ppb glyphosate equivalent, as reflected by changes in function of over 4000 genes, the American public should be concerned about glyphosate residues on their food. Additional research points to harmful impacts at levels between 10 ppb and 700 ppb. Considering these shocking new scientific test results, regulators must take the below findings into account during any re-authorization of glyphosate.



Glyphosate Food Testing Results: (in parts per billion – ppb)

Full laboratory reports for this food testing can be [found here](#). A searchable database of results can be [found here](#).

General Mills		
	Original Cheerios	Glyphosate - 1,125.3 ppb AMPA – 26.4
	Honey Nut Cheerios	Glyphosate – 670.2 ppb AMPA – 14.5
	Wheaties	Glyphosate – 31.2 ppb
	Trix	Glyphosate - 9.9 ppb
	Gluten Free Bunny Cookies Cocoa & Vanilla	Glyphosate – 55.13* ppb
Kellogg's		
	Corn Flakes	Glyphosate – 78.9 ppb
	Raisin Bran	Glyphosate – 82.9 ppb
	Organic Promise**	Glyphosate - 24.9 ppb
	Special K	Glyphosate - 74.6 ppb
	Frosted Flakes	Glyphosate - 72.8 ppb
	Cheez-It (Original)	Glyphosate – 24.6 ppb
	Cheez-It (Whole Grain)	Glyphosate – 36.25* ppb
	Soft-Baked Cookies, Oatmeal Dark Chocolate	Glyphosate – 275.58* ppb
Nabisco		
	Ritz Crackers	Glyphosate – 270.24 ppb
	Triscuit	Glyphosate – 89.68 ppb
	Oreo Original	Glyphosate – 289.47* ppb

Nabisco (continued)		
	Oreo Double Stuf Chocolate Sandwich Cookies	Glyphosate – 140.90* ppb
	Oreo Double Stuf Golden Sandwich Cookies	Glyphosate – 215.40* ppb
PepsiCo		
	Stacy's Simply Naked Pita Chips (Frito-Lay)	Glyphosate – 812.53 ppb
	Lay's: Kettle Cooked Original	Glyphosate – 452.71* ppb
	Doritos: Cool Ranch	Glyphosate – 481.27* ppb
	Fritos (Original) (100% Whole Grain)	Glyphosate - 174.71* ppb
Campbell Soup Company		
	Goldfish crackers original (Pepperidge Farm)	Glyphosate – 18.40 ppb
	Goldfish crackers colors	Glyphosate – 8.02 ppb
	Goldfish crackers Whole Grain	Glyphosate – 24.58 ppb
Little Debbie		
	Oatmeal Creme Pies	Glyphosate – 264.28* ppb
Lucy's		
	Oatmeal Cookies Gluten Free	Glyphosate – 452.44* ppb
Whole Foods		
	365 Organic Golden Round Crackers**	Glyphosate – 119.12* ppb
Back to Nature		
	Crispy Cheddar Crackers	Glyphosate – 327.22* ppb

Limit of Quantitation: 5 ppb

*These samples exhibit very low recovery and/or response. The above amounts found are rough estimates at best and may not represent an accurate representation of the sample.

** Widespread contamination in food supply — even organic farmers are having their crops/ our food contaminated.

The Laboratory

Anresco Laboratories was founded by Dr. Sylvan Eisenberg in 1943, with the goal of ensuring food quality. Their food safety analyses span a variety of areas, including Pesticides, Microbiology and Nutritional Labeling. Since 1980, Anresco has been focused on pesticide and herbicide testing. They perform routine testing on imported samples detained by the FDA, samples to be exported, as well as domestic samples for many major US companies. In addition to foods, they have also tested herbal products, water and soil.

LC/MS/MS is the method used for testing for glyphosate levels in food products. LC/MS/MS stands for liquid chromatography tandem mass spectrometry (LC-MS/MS), which "has been well recognized by regulators and scientists worldwide as the "gold standard" for both quantification and semi-quantitative screening of food contaminants, such as pesticide residues."²⁹

Their instrumentation includes Gas Chromatography (GC-FID, GC-FPD, GC-TSD, GC-ECD, GCMS, GCMSMS) and Liquid Chromatography (LC-FD, LC-UV, LC-RID, LC-MSMS).

The laboratory is an independent, ISO-17025 accredited, FDA-registered private laboratory headquartered in San Francisco, U.S.A.

Anresco Laboratories Co-laboratory Director Vu Lam says, "We use LC/MS/MS to detect Glyphosate and AMPA. With our modifications to various methods, we are able to detect many levels of contaminants in food products at levels as low as 5 ppb, and even lower."

Method for Food Testing

The sample is extracted using water and methanol. Extract is further cleaned and concentrated on an ion-exchange column. The extract is then derivatized using 9-fluorenylmethylchloroformate and filtered prior to LC-MS/MS analysis.

Derivatized glyphosate and its metabolite AMPA are injected onto the LC-MS/MS and separated using a C18 column with a gradient elution of ammonium acetate buffer and acetonitrile as mobile phases. Ionization is achieved using an electrospray ionization source operating in negative mode. Analytes are detected using multiple reaction monitoring with isotopically-labeled analytes as internal standards. This method allows detection and quantification down to 2 ng/g (ppb) for food and soil samples with recoveries between 70-90%.

LC/MS/MS testing is the best method currently available for glyphosate testing in food, soils, urine and water – please see more information on this [here](#).³⁰ Other methods that do not use mass

spectrometry are not recognized by regulators worldwide.

Anca-Maria Tugulea, a chemist in the Exposure and Biomonitoring Division of Health Canada stated:

"ELISA has high false positive and false negative results... It is usually used as a screening tool and any positive results will have to be confirmed by a chromatographic (e.g. LC/MS/MS) method to be usable in risk assessment."³¹



Where Else Has Glyphosate Been Found?

Glyphosate Residues Found in Food, Urine, Breast Milk, Rainwater, Rivers, Tap Water and Tampons – But the FDA Has Never Conducted Proper Widespread Testing

While the U.S. government has never released proper glyphosate residue tests on popular American food products to the public before, this past year on February 17, 2016, the FDA announced that it would finally begin testing glyphosate residues in certain food products "for Fiscal Year 2016 to measure glyphosate in soybeans, corn, milk, and eggs, among other potential foods."³²

Because glyphosate is absorbed into the plant's cellular structure in order to kill weeds or unwanted plants, the chemical cannot be removed from GMO crops and other plants by washing, peeling, baking or brewing grains. In recent years, a growing number of studies have found glyphosate in common foods, human urine,³³ breast milk,³⁴ beer³⁵, 85% of tampons³⁶ and even a group of elected officials in Germany at alarming levels.³⁷

Despite the increased exposure to the American public and citizens around the world, inadequate regulatory testing exists on glyphosate residues in common food products.

According to a U.K. Food Standard Agency test conducted in October 2012, glyphosate residues were found in 27 of 109 samples of bread sold in England at .2 ppm (parts per million).³⁸ While

limited testing conducted by the USDA in 2011 found glyphosate residues on 90.3% of 300 soybean samples tested and 95.7% of AMPA, a metabolite of glyphosate, at concentrations of 1.9 ppm and 2.3 ppm respectively.³⁹

Pure Science: Glyphosate Damage by the Numbers (in parts per billion):

While regulators at the EPA determined in the 1990s that 1.75 mg per kilogram of bodyweight per day is the safe allowable daily intake level in the U.S., the EPA's own website lists glyphosate under "Regulated Drinking Water Contaminants." The EPA warns people that long-term exposure to glyphosate at only 700 ppb can cause "problems with their kidneys or reproductive difficulties" and states that drinking water exposure exists due to "Runoff from herbicide use."⁴⁰

Glyphosate/Roundup Damage by the Numbers (ppb)

0.1 ppb: Roundup (0.05 ppb glyphosate) altered the gene function of over 4,000 genes in the livers and kidneys of rats.⁴¹

0.1 ppb: Roundup (0.05 ppb glyphosate) severe organ damage in rats.⁴²

0.1 ppb: Permitted level for glyphosate and all other herbicides in EU tap water.⁴³

10 ppb: Toxic effects on the livers of fish.

700 ppb: Alterations of kidneys and livers in rats.⁴⁴

700 ppb: Permitted level for glyphosate in U.S. tap water.⁴⁵

1,125.3 ppb (1.1253 mg/kg): Level found in General Mills' Cheerios.

Unfortunately, as a result of the combination of exposures through drinking water and a regular diet of processed food, total exposure is much greater than industry scientists and U.S. regulators ever anticipated. **With this new information, it's time for the EPA to dramatically reduce the acceptable daily intake level of glyphosate and ban the practice of pre-harvest spraying of all food crops, especially wheat, oats and barley, immediately.**



Food Democracy Now! Glyphosate: Unsafe on Any Plate

Alarming Levels of Glyphosate Found Due to Pre-Harvest Spraying

Cheerios

Ingredients: Whole Grain Oats, Corn Starch, Sugar, Salt, Tripotassium Phosphate, Vitamin E (mixed tocopherols) Added to Preserve Freshness.

Vitamins and Minerals: Calcium Carbonate, Iron and Zinc (mineral nutrients), Vitamin C (sodium ascorbate), A B Vitamin (niacinamide), Vitamin B₆ (pyridoxine hydrochloride), Vitamin A (palmitate), Vitamin B₁ (thiamin mononitrate), A B Vitamin (folic acid), Vitamin B₁₂, Vitamin D₃.

DISTRIBUTED BY GENERAL MILLS SALES, INC.,
MINNEAPOLIS, MN 55440 USA

GLUTEN FREE

Not made with genetically modified ingredients.

Trace amounts of genetically modified (also known as "genetically engineered") material may be present due to potential cross contact during manufacturing and shipping.

Non-GMO Cheerios (Yellow box) 1,125.3 ppb



INGREDIENTS: Enriched Wheat Flour (Wheat Flour, Niacin, Reduced Iron, Thiamin Mononitrate, Riboflavin, Folic Acid), Sunflower Oil and/or Canola Oil, Sea Salt, Whole Wheat Flour, and Less than 2% of the Following: Organic Cane Sugar, Oat Fiber, Yeast, Malted Barley Flour, Rosemary Extract (Antioxidant), and Ascorbic Acid (Antioxidant).

CONTAINS WHEAT INGREDIENTS.

Stacy's Pita Chips (Simply Naked) 812.53 ppb



Less Than 0.9% GMO,
But Contains Glyphosate

What Do These Numbers Mean?

To understand what the above numbers mean to the average consumer, consider the fact that the acceptable daily intake (ADI) level is currently set at 1.75 milligrams per kilogram of bodyweight per day by the EPA. This means that over the course of a day, an individual would have to combine all the residue levels found on their meals and snacks and consistently be below this supposedly "safe" level.

According to the old theory "the dose makes the poison", any individual whose daily exposure is at or below this level over the course of their lifetime should not have any negative health impacts. The current ADI level was determined using industry funded studies from the 1970s, and 1980s and according to the latest advances in scientific research is woefully out of date.

Harmful Effects of Chemicals at Low Levels and the Flawed Logic Behind "the Dose Makes the Poison"

Revelations from the latest scientific research finds that chemicals can have damaging effects at very low levels. In fact, since Roundup and glyphosate were originally approved and declared "safe", a whole new body of research finds that chemicals with endocrine (or hormone) disrupting capabilities, turns the "dose makes the poison" logic on its head, finding that low level doses or exposures to certain chemicals can have much larger negative impacts on health and human development than previously understood.

For more than two decades, research has shown that low level exposures to certain chemicals, at levels previously considered safe, can alter the release and reception of vital hormones in the body.

Significant scientific evidence points to the fact that low levels exposure to endocrine disrupting chemicals can cause reproductive problems, reduced fertility, miscarriage and that changes in hormone levels can result in the early onset of puberty, obesity, diabetes, behavioral problems, including attention deficit/hyperactivity disorder (ADHD), impaired immune function and certain types of cancers.†

And today, while glyphosate has not been definitively declared an endocrine disruptor (or hormone hacker), new and emerging evidence suggests that it may be and independent scientists are finding an increasing number of significant

harms from glyphosate at very low levels previously declared "safe" by the EPA.

Our current regulatory system relies on outdated notions of risk and exposure by relying on safety assessments that only call for high-dose exposure studies, which are used to determine "safe" exposure levels. With the emergence of science proving endocrine disrupting chemicals are pervasive in our food and environment, critics of the current system signed a consensus statement that "The very low-dose effects of endocrine disruptors cannot be predicted from high-dose studies, which contradicts the standard 'dose makes the poison' rule of toxicology."‡

Real World Glyphosate Exposure at Detected Levels

To calculate real world exposure levels from glyphosate contamination in our food, consider the Cheerios' results, since they're the highest and are also regularly consumed by children, a chemically vulnerable population.

For a 1-year old child, at roughly 20 pounds (9 kg), General Mills recommends a serving size of 16 grams or .56 ounces.

Based on OLD SCIENCE: Acceptable Daily Intake (ADI) set by government regulators: 1.75 milligrams / kilogram of bodyweight / per day (1.75 mg / kg bw / day)

Daily Intake from Cheerios (DI) based on 1 serving of 16 grams containing 0.018 mg glyphosate equals 0.002 mg / kg bw/day for a 20lb (9kg) child.

So, for a 20-pound child:

NEW SCIENCE says that harm may happen when you eat glyphosate at ONLY 0.001 mg / kg bw/day (1,750 times LOWER than what the EPA at the moment says is safe.) So, New Science shows that a 20lb (9kg) child can ONLY safely eat 0.001mg x 9kg = 0.009mg glyphosate in a day.

Thus, ingesting a recommended serving of Cheerios (16 grams or .56 ounces) for a 20-pound (9kg) child means exposure to 0.018 mg of glyphosate per day from Cheerios alone. This is nearly double the level that is potentially harmful to human health and it is important to note that a 1-year old child is likely to be exposed to many other sources of glyphosate on a daily basis.

† Fürst P (October 2006). "Dioxins, polychlorinated biphenyls and other organohalogen compounds in human milk. Levels, correlations, trends and exposure through breastfeeding". *Mol Nutr Food Res.* 50 (10): 922-33. doi:10.1002/mnfr.200600008. PMID 17009213.

‡ Collaborative on Health; the Environment's Learning; Developmental Disabilities Initiative (2008-07-01). "Scientific Consensus Statement on Environmental Agents Associated with Neurodevelopmental Disorders", Institute for Children's Environmental Health. Retrieved 2009-03-14.

Reasons for Concern: Glyphosate Risks to Human Health

1. Cancer

Despite repeated claims from Monsanto that glyphosate is perfectly safe, even “safer than table salt,” an increasing number of independent peer-reviewed studies are finding troubling links to a growing number of diseases in humans, including cancer.

In March 2015, leading cancer experts from the World Health Organization’s International Agency for Research on Cancer (IARC) declared that glyphosate is a “probable human carcinogen.” IARC reached its decision based on the research of 17 top cancer experts from 11 countries, who met to assess the carcinogenicity of 5 pesticides. The IARC review of glyphosate has led to the European Parliament calling for a complete ban on non-commercial public use of glyphosate and serious restrictions on agricultural use.⁴⁶

Already, multiple class action lawsuits⁴⁷ and a wrongful death lawsuit⁴⁸ have been filed in the U.S. against Monsanto, based on the finding that glyphosate is a probable cancer-causing agent used with regularity by farmers, farm workers, pesticide applicators and others exposed to the chemical through routine use. There are specific concerns⁴⁹ over the following cancers:

- Non-Hodgkin’s Lymphoma
- Bone Cancer
- Colon Cancer
- Kidney Cancer
- Liver Cancer
- Melanoma
- Pancreatic Cancer
- Thyroid Cancer

For More Information on Glyphosate and Cancer: <http://detoxproject.org/glyphosate/cancer/>

2. Endocrine Disruption and Hormone Hacking Capabilities

A growing body of peer-reviewed scientific evidence shows that glyphosate also acts as an endocrine disruptor at levels that the human population is routinely exposed to in America through the consumption of unlabeled GMO foods and foods that have been sprayed with glyphosate as a pre-harvest drying agent.

Incredibly, the potential harm to human reproductive capability as a result of glyphosate’s toxicity has never been studied by regulators or the chemical industry at levels that the human population in the U.S. is being exposed to (under 3 mg/kg body weight/day) on a daily basis. This is a huge hole in the global risk assessment of glyphosate, as there is evidence suggesting that low levels of some chemicals may hack hormones more than at mid and high levels, according to independent science – in other words, a higher dose does not necessarily make a more toxic, hormone disruptive effect.

A study from March 2015 stated that the health costs to the European Union of just some endocrine disrupting hormone-hacking chemicals, in connection with a subset of illnesses known to be linked to hormone interference, is over €150 billion (Euros) per year.⁵⁰ The study stated that lower IQ, adult obesity and 5% or more of autism cases are all linked to exposure to endocrine disruptors.

For More Information on Glyphosate and Endocrine Disruption or Hormone Hacking: <http://detoxproject.org/glyphosate/hormone-hacking/>

3. Binding of vital nutrients

Glyphosate binds (chelates) vital nutrients⁵¹ such as iron, manganese, zinc, and boron in the soil, preventing plants from taking them up, which could mean loss of basic minerals and nutrients in food crops and soil as a result. This has implications for humans, livestock and pets that eat glyphosate-tolerant GMO crops, as it affects nutritional value.

For More Information on How Glyphosate Binds with Vital Nutrients: <http://detoxproject.org/glyphosate/glyphosate-chelating-agent/>

4. Antibiotic or Biocidal Properties Cause Concern

Glyphosate is also patented as an antibiotic or biocide, meaning that beyond its use as an herbicide to kill weeds it in most likelihood has a significant harmful effect on humans and farm animals by killing beneficial microorganisms in the gut. In 2010 Monsanto was awarded a patent by the U.S. patent office, which it originally applied for in 2003, in order to use glyphosate to kill “protozoan parasites.” In their patent application Monsanto is very clear about the chemical’s ability to kill parasites.

From Monsanto’s U.S. Patent #7,771,736 Glyphosate formulations and their use for the inhibition of 5-enolpyruvylshikimate-3-phosphate synthase:

“Importantly, the growth of these parasites can be inhibited by the herbicide glyphosate, suggesting that the shikimate pathway will make a good target for the development of new anti-

parasite agents. The present invention discloses the use of the herbicidal agent glyphosate in combination with the polyvalent anion oxalic acid for the prevention and therapy of these pathogenic infections.”

While Monsanto continues to deny any possible harmful health impacts, new research points to growing concerns from independent scientists and medical professionals regarding glyphosate’s likely antimicrobial effects on human and animal health. Evidence suggests that glyphosate can alter the balance of the gastrointestinal microbiome, or beneficial gut bacteria, and create a harmful imbalance in human and animal gut flora while increasing pathogenic species that can lead to multiple negative health outcomes, including gluten-intolerance and irritable bowel syndrome.

In the first such research in the world, a team from the University of Canterbury in New Zealand found in 2015 that commonly used herbicides, including Roundup (glyphosate-based), can cause bacteria to become resistant to antibiotics.⁵²

Not surprisingly, this study was picked up by U.S. cardiologist Dr. William Davies, *the New York Times* bestselling author of *Wheat Belly*, which details the alarming rise in gluten sensitivity and intolerance in the American public, who noted the study’s details on his website:

“Animal model data demonstrates that glyphosate selectively kills beneficial bacteria, such as *Enterococcus faecalis*, *Enterococcus faecium*, *Bacillus badius*, *Bifidobacterium adolescentis* and *Lactobacillus* species, while allowing the proliferation of undesirable, even disease-causing, species such as *Salmonella*


enteritidis, *Salmonella gallinarum*, *Salmonella typhimurium*, *Clostridium perfringens* and *Clostridium botulinum*.”⁵³

In 2012, a team of German scientists investigated the rise in botulism disease in cattle in the past 10–15 years found that glyphosate and Roundup were toxic to beneficial gut bacteria that inhibit the growth of the botulism-causing bacterium *Clostridium botulinum*, but non-toxic to the botulism-causing bacteria themselves. In short, glyphosate and Roundup favored the growth of botulism-causing *Clostridium botulinum* bacteria. The authors concluded that ingestion of Roundup residues in cattle feed could predispose cattle to falling ill with botulism.⁵⁴

In a separate in vitro study on strains of bacteria found in the gut of poultry, most of the pathogenic bacteria tested were highly resistant to Roundup, but most of the beneficial gut bacteria tested were found to be moderately to highly susceptible. The researchers documented the antibiotic damage done to beneficial bacteria in the gut by very low concentrations of Roundup, which allowed the overgrowth of serious pathogens such as *Clostridium botulinum*, *Salmonella* spp, and *E. coli*. These would otherwise be kept in check by the beneficial bacteria that were wiped out by the Roundup residues in feed.⁵⁵

The authors concluded that the ingestion of Roundup-contaminated feed could be a significant factor predisposing poultry to diseases caused by *Clostridium botulinum*. It could also explain the now widespread contamination of poultry products with pathogenic *Salmonella* and *E. coli* strains of bacteria, which can make human consumers ill.

Peer-Reviewed Studies on Low Doses of Glyphosate / Roundup



Source: Mesnage, 2015 <http://dx.doi.org/10.1016/j.fct.2015.08.012>

Food Democracy Now! Glyphosate: Unsafe on Any Plate

Antibiotic resistance is a serious and growing problem for human and animal health. New antibiotics are hard to find and can take decades to become available.

Further studies are now needed to see how glyphosate affects the gut microbiome in humans and to see if glyphosate at the actual doses ingested is a cause of antibiotic resistance in humans.

Where Do Claims of “Safe” Glyphosate Levels Come From?

Industry and regulators have often claimed that harmful effects caused by glyphosate and Roundup in animal studies do not matter because the public is only exposed to “safe” levels that do not cause such effects.

Currently, regulators in the U.S. and Europe only set safety limits for exposure to glyphosate based on data from industry’s own toxicity studies on laboratory animals. The results of these industry tests, which are conveniently classified as commercial secrets and kept hidden from the public and the independent scientific community, are presented to expert panels in government or food safety agencies based in various countries or regions.

Unfortunately, the U.S. EPA and the European Food Safety Authority (EFSA) refuse to release the list of all the chemical ingredients used in the most common pesticide mixes, once again siding with industry lobbyists and hiding these toxic chemicals from the American public, all in the name of “trade secrets.”

Reasons for Concern: Allowed Levels of Glyphosate Are Unlikely to be Safe

There are several reasons to doubt the validity of the current Acceptable Daily Intake (ADI) levels for glyphosate, including:

- The so-called safe levels of glyphosate exposure have never been tested directly to determine if indeed they are really safe to consume over the long term. Instead the “safe” levels are extrapolated from higher doses tested in industry studies.
- Industry toxicity study protocols are massively out of date. All toxicity tests conducted by industry for regulatory purposes are based on the old adage, “The dose makes the poison,” – that is, the higher the dose, the greater the

degree of toxicity. However, in some cases, low doses corresponding to human exposures can be more toxic than the higher doses tested in laboratory animals in industry studies. This is especially true for chemicals that disrupt the hormonal system (endocrine disruptors).⁵⁶

- Safe levels of these endocrine disrupting chemicals cannot be extrapolated from effects at higher doses. Evidence from in vitro⁵⁷ and animal⁵⁸ experiments shows that glyphosate may be an endocrine disruptor at levels permitted in tap water in the EU and the U.S.
- Findings that glyphosate and its commercial formulations may be endocrine disruptors imply that the standard industry long-term animal studies are seriously inadequate.
- Hormones are vital regulators of fetal development. A subtle hormonal effect during early life can modify organ morphology and function for the rest of the life, as well as potentially leading to chronic diseases such as cancer and neurological and reproductive dysfunction in adults⁵⁹.
- Even worse, the complete glyphosate herbicide formulations as sold and used contain additives (adjuvants), which are more harmful in their own right and/or increase the toxicity of glyphosate.⁶⁰ Safety limits are set for the isolated ingredient glyphosate, but the whole formulations, which can be generally more toxic, are never tested to determine long-term effects. This limitation of the regulatory process applies to all pesticides in all countries worldwide. Studies in rats confirm that the complete glyphosate herbicide formulations are toxic at levels deemed by regulators to be safe or have no adverse effect for the isolated ingredient glyphosate.^{61,62,63} Other feeding studies in pigs⁶⁴ and rats⁶⁵ directly comparing the toxicity of formulations with glyphosate alone found that the formulations were far more toxic.
- Industry tests on glyphosate alone revealed toxic effects, notably birth defects^{66,67}, below the levels that regulators claimed showed no toxic effect – but these results were ignored or dismissed by regulators in setting the supposedly safe ADI.⁶⁸
- Independent studies have found toxic effects of glyphosate and its commercial formulations at environmentally realistic levels, which have never been tested by regulators. Effects include oxidative stress on liver and kidneys⁶⁹ and endocrine disrupting effects.⁷⁰
- Glyphosate, which was claimed to be “as safe as table salt”⁷¹ by Monsanto for over 40 years, was classified as a probable human

carcinogen by the World Health Organization in 2015. Glyphosate has never been tested during sensitive periods of life (such as fetal development) at environmentally relevant levels of exposure. In addition, the fact that Monsanto's commercial formulations have never been tested for regulatory purposes for more than one month in rats, and that without any blood testing, raises further doubt as to the validity of current ADI values.

These findings, taken as a whole, suggest that the levels of Roundup or glyphosate the American public are exposed to are not safe over the long term.

What Have U.S. Food Safety Regulators Done to Protect Us?

Obama Administration Raised Glyphosate Food Residue Levels at Monsanto's Request

Rather than call for increased oversight and monitoring of glyphosate in popular American foods, in 2013, the U.S. Environmental Protection Agency, under the direction of the Obama administration, actually increased the allowed tolerance pesticide exposure levels on glyphosate residues based on a petition to the agency submitted by Monsanto.^{72 73 74}

These latest increases in glyphosate food residues in the U.S. raised allowable levels in oilseed crops, which include sesame, flax, and soybean, from 20 parts per million (ppm), to 40 ppm.

Under the Obama administration, the EPA also increased allowable glyphosate contamination levels for sweet potatoes and carrots from 0.2

ppm to 3 ppm for sweet potatoes and 5 ppm for carrots. That's 15 and 25 times the previous levels, respectively.⁷⁵

An independent, peer-reviewed study conducted in 2014 on Roundup Ready soybeans found "extreme levels" of glyphosate residue in 7 out of the 10 GMO soy samples tested when compared to organic and conventionally raised soybeans.^{76 77}

If these facts weren't troubling enough, recent testing commissioned by the Organic Consumers Association in coordination with The Detox Project at the University of California San Francisco (UCSF), found glyphosate in 93% of Americans tested at an average level of 3.096 parts per billion (ppb). Children had the highest levels with an average of 3.586 ppb.^{78 79}

Acceptable Daily Intake (ADI) or How Much Pesticide Residue You May Eat Every Day

Real concerns about glyphosate residue levels are growing in the U.S. and around the world, not only due to Monsanto's long-standing reputation for producing toxic chemicals that cause severe harm to human health and the environment, e.g. Agent Orange, DDT and PCBs, but also because of the persistence of glyphosate in food crops, which this study helps to confirm.

A 2014 study found that high levels of glyphosate exposure in animals and humans were correlated with an increase in chronic illness. The authors discovered that:


"Glyphosate residues in urine were grouped according to the human health status. Chronically ill humans had significantly higher glyphosate residues in urine than healthy humans."⁸⁰

In order to limit threats to human health, federal regulatory agencies set limits on acceptable exposures based on a safety threshold, also called the Acceptable Daily Intake (ADI).⁸¹ To assess whether or not an exposure is a health risk, the ADI or safety threshold for a given chemical is an estimate of the amount of a substance in food or drinking water, expressed on a body mass basis, that can be ingested daily over a lifetime by humans without appreciable health risks.

This calculation of Allowable Daily Intake for pesticide residues is based on the lowest dose considered non-toxic in animal feeding trials (i.e. 30 mg/kg bw/d) sponsored by industry.

Accordingly, based on Monsanto's own private lab testing and consulting with scientists at different regulatory agencies around the world, the ADI value for glyphosate differs from country to country.

U.S. Government Allows Nearly 6 Times More Glyphosate in Our Food than Europe



In Europe, the current ADI has been set at 0.3 mg per kg of bodyweight per day (written as 0.3 mg/kg bw/d),⁸² while in the United States glyphosate's allowable daily intake is nearly 6 times higher at 1.75mg/kg bw/d.

These wide differences in allowable daily residue exposure mean that U.S. citizens are legally exposed to nearly 6 times the amount of glyphosate on a daily basis than individuals in Europe.

Changes in Daily Exposure Based on Industry Science, at Monsanto's Request and a History of Scientific Fraud

In the case of glyphosate, this Acceptable Daily Intake level has been based on limited scientific studies presented to the EPA by glyphosate's original applicant for approval, Monsanto. As in Europe, the U.S. EPA has historically made these decisions based on corporate in-house scientific studies that have never gone through the peer review process, nor has any of the original data from these studies reported to regulatory agencies ever been made public.

More importantly, the differences in current Acceptable Daily Intake levels between the European Union and the U.S. are significant, do not represent the best or most current scientific data and are shrouded in controversy. This is not to say that this report endorses the European Union standard, which recent scientific evidence suggests establishes a tolerance too high to be guaranteed to safeguard human health.

U.S. Acceptable Daily Intake for Glyphosate (ADI) Originally Set by EPA at 0.1 ppm

According to internal EPA documents, the original ADI set by the EPA toxicology division was 0.1 mg/kg/day in the early 1980s.⁸³ During this same period under the Reagan Administration, in 1985, the EPA classified glyphosate as a possible carcinogen, Class C, based on a long-term feeding study in which male mice fed glyphosate developed kidney tumors.⁸⁴

The EPA initially defended this position, but Monsanto successfully submitted "historical control data" from multiple other unpublished studies and in a June 26, 1991 memo, the EPA reclassified glyphosate as Class E or "non-carcinogenic for humans" after much back-and-forth for several years with Monsanto scientists and lobbyists "based upon lack of convincing carcinogenicity evidence in adequate studies in two animal species."⁸⁵

While Monsanto and other chemical manufacturers in the United States defend the low toxicity of Roundup and other glyphosate-based herbicides,

regulators in Europe took a more cautious scientific approach in setting the Acceptable Daily Intake limits for their citizens.

Rather than take Monsanto's approach on allowable levels, in its 1998 evaluation of glyphosate, Germany's Federal Office of Consumer Protection and Food Safety (BVL) determined that the allowable residue level should be set at 0.3 mg per kilogram of body weight, versus the high level of 1.75 ppm set by the EPA, or nearly 6 times higher than acceptable levels allowed in Europe based on industry feeding trials that they believed to be the most sensitive to the effects of the chemical.⁸⁶

The decision-making process of the German government's Consumer Protection and Food Safety agency was spelled out in a public document that stated obvious concern over the high ADI chosen by their U.S. counterparts at the EPA. According to the BVL glyphosate review:

"A very high ADI of 1.75 mg/kg bw was proposed in the joint dossier of Monsanto and Cheminova and is based on the NOEL for maternal toxicity in a teratogenicity study in rabbits (Tasker, 1980). It is discussed here since it is far outside the range of all the other suggested values."⁸⁷

The German review document details the then-current ADI level requests by various chemical manufacturers based on industry-approved studies made available to respective food safety agencies around the world in the 1990s. These original Acceptable Daily Intake levels range from 0.05 to 0.1 mg/kg bw/day, 0.15 mg/kg bw/day and 0.3 mg/kg bw/day and the requested U.S. level of 1.75 mg/kg bw/day submitted by Monsanto and Danish pesticide maker Cheminova. (See original chart on page 16)

It's interesting to note that the U.S. ADI level of 1.75 mg/kg bw/day is 17.5 times the original ADI set by the EPA in the 1980s⁸⁸ and was obviously raised in anticipation of the approval of future GMO crops. Monsanto had already begun advance work on engineering genetically modified crops that were design to survive being sprayed with their proprietary flagship herbicide Roundup.

In their 1998 review of glyphosate, German regulators stated their objections to the EPA and Monsanto's ADI request in clear scientific terms:

"The acceptable daily intake should be based on the highest dose at which no adverse effect is observed in the most appropriate study in the most sensitive species. In the case of glyphosate, the different notifiers have proposed ADI values which cover a wide range between 0.05 and 1.75 mg/kg bw (see table B.5.10.2-1). This variance is due to the different studies used as

the respective basis for ADI calculation but may also result from a controversial evaluation of controversial studies.

A very high ADI of 1.75 mg/kg bw was proposed in the joint dossier of Monsanto and Cheminova and is based on the NOEL for maternal toxicity in a teratogenicity study in rabbits (Tasker, 1980). It is discussed here since it is far outside the range of all the other suggested values. This proposal was not accepted by the Rapporteur for the following reasons:

1. The NOEL for maternal toxicity in the respective study was established by the Rapporteur at 75 mg/kg bw/day instead of 175 mg/kg bw/day (see section B.5.6.2.2.2).
2. If a NOEL of 175 mg/kg /bw/day for the above mentioned rabbit study would have been accepted, one could identify some valid studies revealing adverse effects at lower doses. In a recent long-term study in rats (Suresh, 1996), effects occurred in female animals at a dietary dose level of 1000 ppm (ca 60/mg/bg bw/day). **The NOELs [No Observed Effect Level] and LOELs [Lowest Observed Effect Level] established in a further chronic rat study (Atkinson et al., 1993) and in two other rabbit teratogenicity studies (Suresh, 1993; Brooker et al., 1991) were well below 175 mg/kg bw/day.**


Usually, a chronic rat study is considered most appropriate to derive the ADI. Since the rat proved the most sensitive species upon long-term exposure, it is suggested to establish the ADI for glyphosate on the basis of the chronic toxicity data obtained in rats.⁶⁹

Current Scientific Research Calls for Much Lower Allowable Glyphosate Residues for Human Food Products

This spring, as regulators in the European Union attempted to re-register glyphosate for another 15 years, new independent scientific evidence was brought forward that called into question even the EUs more cautious 0.3 mg/kg bw/day allowable glyphosate residue level.

In the past 10 years alone, real scientific concern over the chemical's safety has only increased due to the widespread explosion in the use of Roundup and glyphosate-based herbicides in conjunction with industrial agriculture and further pressures on farmers to adopt GMO agriculture linked to herbicide tolerant traits.

In a study titled "*Transcriptome profile analysis reflects rat liver and kidney damage following chronic ultra-low dose Roundup exposure*," a team of international scientists performed a



transcriptome (gene expression) analysis of the liver and kidneys from rats fed an extremely low dose of Roundup. This resulted in a daily intake of glyphosate of only 4 nanograms per kilogram of bodyweight per day, which is 75,000 times below the EU acceptable daily intake (ADI equivalent) and 475,000 times below the US chronic reference dose (ADI equivalent). In other words, **a dose of Roundup that was far below what is permitted by regulators and believed to be safe to consume on a daily basis over the long term was found to damage the liver and kidneys of rats.** These results were statistically significant.⁹⁰

While transcriptome analysis is highly predictive of disease status or organs, it cannot predict the corresponding disease states with absolute certainty, as not all changes in gene function result in changes in levels of the genes' protein products and metabolites. Such definitive proof has to be provided by additional molecular profiling analysis, namely proteomics (protein profile) and metabolomics (small molecule metabolite profile). The proteomics and metabolomics analyses give a direct measure of the organ's composition, so they are able to provide a direct indicator of the health or disease status of the organ in question.

The authors concluded that long-term exposure to Roundup “at an ultra-low, environmental dose can result in liver and kidney damage with potential significant health implications for animal and human populations.”

—MESNAGE ET AL, 2015

A separate study looked at the evidence for teratogenic effects (birth defects) in the industry studies submitted to regulators to gain market authorization for glyphosate. The authors reviewed

the German government’s original 1998 scientific analysis of allegedly safe daily glyphosate exposure levels and found that the government regulators had “introduced significant bias” into their review by intentionally excluding toxicity studies in rabbits that found harmful effects of glyphosate at much lower levels than their analysis admitted.^{91,92}

Simply by reviewing the same studies that the German government regulators did in 1998, the authors calculated that the EU acceptable daily intake level of glyphosate was currently 3 times higher than it should be if all the industry studies had been rigorously evaluated.

By including in their analysis two independent peer-reviewed studies that had been completed since the EU set the acceptable daily intake for glyphosate residue in 2002, **the authors concluded that current ADI should be dramatically reduced to 0.025 mg/kg bodyweight per day or “12 times lower than the ADI proposed by the German regulators, which is currently in force in the EU and used as a basis for the maximum residue limit for food and feed.”**

Multi-Generational Rat Studies on Glyphosate with Recommended ADI Levels

Table B.5.10.2-1: Summary of ADI values proposed by the different notifiers and by the Rapporteur

Notifier	ADI (mg/kg bw)	Toxicological data on which this ADI proposal is based	Remarks of the Rapporteur
Monsanto/ Cheminova	1.75	Teratogenicity study in rabbits, NOEL: 175 mg/kg bw/d.	See discussion below.
Agrichem	0.1	3-generation study in rats, NOEL 10 mg/kg bw/d.	Based on published literature. Study not identified. Much higher NOELs have been established in more recent reproduction studies.
Alkaloida	0.06	12-month study in dogs, NOAEL: 300 ppm (5.79 - 14.62 mg/kg bw/d).	Supplementary study, NOAEL = highest dose tested.
Barclay	0.3	Chronic study in rats (NOEL 31 mg/kg bw/d) and 3-generation study in rats (NOEL 30 mg/kg bw/d) with reference to 1986 JMPR evaluation.	No original studies. In both cases, the NOELs were the highest doses tested. Both studies were considered supplementary only.
Feinchemie	0.05	Chronic study in rats, NOEL: 100 ppm (ca 5.5 mg/kg bw/d).	Interim report conclusion.
Herbex	-	Proposal for an ADI not submitted; appropriate studies not performed.	-
Luxan	0.15	Carcinogenicity study in mice (NOAEL 150 ppm, ca 15 mg/kg bw/d) and 3-generation study in rats (NOEL 300 ppm, ca 15 mg/kg bw/d).	Supplementary studies. In the reproduction study, NOEL = highest dose tested. Much higher NOELs have been established in more recent long-term and reproduction studies.
Nufarm	-	No toxicological data submitted.	-
Sanachem	0.3	Published literature.	It is assumed that this value refers to the JMPR evaluation in 1986 (i.e. ADI derived from a long-term rat study).
SCC/I.Pi.Ci.	-	Proposal for an ADI not submitted; appropriate studies not performed.	The company refers to the database of other notifiers.
Sinon (Shinung)	0.3	Published literature.	It is assumed that this value refers to the JMPR evaluation in 1986 (i.e. ADI derived from a long-term rat study).
Rapporteur	0.3	Summary of long-term studies in rats.	See discussion below.

Source: BVL, Germany, 1998. Monograph on Glyphosate.

Food Democracy Now! Glyphosate: Unsafe on Any Plate

Monsanto and EPA Claim Roundup and Glyphosate Are Perfectly Safe: Science Says Otherwise

Roundup Formula 125 Times More Toxic than Glyphosate Alone

For decades, Monsanto has publicly claimed that glyphosate was perfectly safe and the company's Roundup herbicide formulations are: "tough on plants, but no more toxic to people and animals than table salt" or "practically non-toxic."⁹³

However, an increasing number of independent peer-reviewed studies have proven that glyphosate is not the most toxic ingredient in Monsanto's Roundup formulation.^{94 95} Glyphosate is never applied to farmers' fields by itself. Instead, pesticide manufacturers create chemical formulations with added ingredients, called "inerts" or "adjuvants," that are needed to penetrate the plant's cell walls to deliver glyphosate into the plant's growth structure. There it works to block the synthesis of three aromatic amino acids essential for growth and makes the plant susceptible to disease and "soil borne fungal pathogens."^{96 97}

According to an independent peer reviewed study published in the International Journal of Environmental Research and Public Health in 2014, scientists found that Roundup was 125 times more toxic than glyphosate alone:

"It is commonly believed that Roundup is among the safest pesticides. This idea is spread by manufacturers, mostly in the reviews they promote... which are often cited in toxicological evaluations of glyphosate-based herbicides. However, Roundup was found in this experiment to be 125 times more toxic than glyphosate. Moreover, despite its reputation, Roundup was by far the most toxic among the herbicides and insecticides tested. This inconsistency between scientific fact and industrial claim may be attributed to huge economic interests, which have been found to falsify health risk assessments and delay health policy decisions."⁹⁸

As a result of new and emerging research, several European countries have not only banned Roundup and glyphosate for use in public parks or sale for home gardens, but the German and French governmental health and safety agencies have forced pesticide manufacturers to remove at least one inert ingredient or "co-formulant" from Monsanto's Roundup Classic and Roundup Original formulas.⁹⁹

For decades, the dangerous chemical known as polyethoxylated tallow amine (POEA), which is derived from animal fat, was a central ingredient in Monsanto's Roundup formula, making up to 15% of

the Roundup Original chemical mixture. As early as 2009 the prestigious Scientific American magazine reported that research scientists had found that:

POEA was more deadly to human embryonic, placental and umbilical cord cells than the herbicide itself – a finding the researchers call "astonishing."¹⁰⁰

According to the 2009 study, published in Chemical Research in Toxicology, "Moreover, the proprietary mixtures available on the market could cause cell damage and even death [at the] residual levels" found on Roundup-treated crops, such as soybeans, alfalfa, corn, and lawns and gardens.¹⁰¹

Scientific American further reported concerns from a team of research scientists, who "suspects that Roundup might cause pregnancy problems by interfering with hormone production, possibly leading to abnormal fetal development, low birth weights or miscarriages."

Despite these concerns and the mounting scientific evidence of likely harm from Monsanto's Roundup formulas, regulators at the USDA and EPA have failed to incorporate this new research into their consideration of Roundup's potential toxicity.

Glyphosate Bio-Accumulates in Major Organs and Bones

While Monsanto and U.S. regulatory agencies routinely claim that glyphosate is excreted quickly from the body, a number of studies in Europe have discovered higher levels of glyphosate residue found in cows raised in countries where GMO feed was allowed (Denmark) and significantly lower in areas considered "GM free" (Germany).¹⁰²

Despite Monsanto's repeated claim that glyphosate does not bio-accumulate,¹⁰³ this 2014 study found glyphosate residues in multiple organs of slaughtered cows, including the intestine, liver, muscles, kidney and spleen, bringing into question Monsanto's claim that glyphosate is rapidly excreted and does not bio-accumulate in animals or humans.

Beyond accumulation in vital organs, glyphosate has also been found to accumulate in bones due to its strong chelating activity or ability to bind with calcium. According to the EPA's own internal documents, reporting on corporate-paid studies submitted by Monsanto, a significant portion of glyphosate is absorbed into the bones of mice and rats used in laboratory experiments.

In 1993, in the EPA's Reregistration Eligibility Decision (RED) on Glyphosate as reported by the Office of Prevention, Pesticides and Toxic Substances: "Less than 10% of the absorbed dose remained in tissues and organs, primarily in bone tissue."¹⁰⁴ The real question remains, what impact

does this steady accumulation in bone tissue have on human health?

In a study on humans and livestock, scientists found that "chronically ill humans had significantly higher glyphosate residues in urine than healthy humans" and also discovered that humans who ate conventional diets had much higher glyphosate residues than those who ate organic food.

According to the 2014 peer reviewed study published in the Journal of Environmental & Analytical Toxicology:

"Glyphosate was significantly higher in humans [fed] conventional [food] compared with predominantly organic [fed] humans. Also the glyphosate residues in urine were grouped according to the human health status. Chronically ill humans had significantly higher glyphosate residues in urine than healthy humans."¹⁰⁵

If these new findings weren't disturbing enough, a series of recent independent peer-reviewed studies found that low doses of Roundup or glyphosate-based herbicides were likely to damage the liver and kidneys of rats at ultra-low dose levels "in the range of what are now generally considered 'safe' for humans."¹⁰⁶

New scientific evidence of the harm from glyphosate continues to emerge at an almost dizzying pace, with concern in the scientific community linking Monsanto's most widely used weedkiller to endocrine disruption, disturbance of normal gut bacteria, autoimmune diseases, birth defects, reproductive problems, infertility and even potential antibiotic resistance.

New research is finding that some autoimmune and chronic inflammatory disorders such as rheumatoid arthritis are associated¹⁰⁷ in other studies with an increased risk of certain types of cancer, including non-Hodgkin's lymphoma, which was correlated with glyphosate exposure in the assessment by the World Health Organization's cancer agency IARC.¹⁰⁸ These findings raise the possibility that rheumatoid arthritis and non-Hodgkin's lymphoma share a common causative factor.^{109 110}

For more than four decades, Monsanto has claimed that glyphosate did not bioaccumulate in animals or humans in any significant way, but a review of a 2004 joint report on pesticide residues in food by the United Nations Food and Agriculture Program and the World Health Organization determined that glyphosate does accumulate in the bones of lab animals.

"Analysis of individual tissues demonstrated that bone contained the highest concentration of [¹⁴C] glyphosate equivalents (0.3–31ppm). The remaining tissues contained glyphosate

equivalents at a concentration of between 0.0003 and 11ppm (Table 3). In the bone and some highly perfused tissues, levels were statistically higher in males than in females."¹¹¹

The question is, since non-Hodgkin's lymphoma is a cancer that starts in cells called lymphocytes, which are part of the body's immune system and can be found in bone marrow, what impact does this daily exposure to increasing levels of glyphosate residues have on the American public, which relies on a diet of processed foods, more than 75 percent of which contain genetically engineered ingredients that were sprayed with high levels of Roundup, Monsanto's flagship weedkiller?

Despite Monsanto's repeated claims of Roundup's safety, the company was successfully sued by the New York state's attorney general in 1996 over its use of "false and misleading advertising," which forced the company to stop claiming its weedkiller was "biodegradable" and to halt ads that claimed Roundup was "safer than table salt" and "practically non-toxic".¹¹²

A similar lawsuit was filed in France that resulted in a former chairman of Monsanto Agriculture France being "found guilty of false advertising for presenting Roundup as biodegradable and claiming that it left the soil clean after use" and a small fine for Monsanto's French distributor.¹¹³

Peer-Reviewed Science on Glyphosate

There are many independent peer-reviewed studies that show the damage caused by glyphosate to human, animal and environmental health. Many of these studies can be found [here](#).

Some of the most important studies and reviews are summarized below:

Cancer

- The teratogenic potential of the herbicide glyphosate-Roundup in Wistar rats. <https://www.ncbi.nlm.nih.gov/pubmed/12765238>
- Two long-term studies on rats indicating possible carcinogenic effects already existed at this time. These long-term studies on rats were conducted in 1979–1981 and 1988–1990. <http://www.inchem.org/documents/ehc/ehc/ehc159.htm>

Human Epidemiological Studies Confirming Cancer Risk

Studies in human populations have found an association between Roundup exposure and two types of blood cancer:

- An epidemiological study of pesticide



applicators in the USA found that exposure to glyphosate herbicide was associated with higher incidence of multiple myeloma. <http://www.ncbi.nlm.nih.gov/pubmed/15626647>

- Epidemiological studies conducted in Sweden found that exposure to glyphosate herbicide was linked with a higher incidence of non-Hodgkin's lymphoma. <http://onlinelibrary.wiley.com/doi/10.1002/%28SICI%291097-0142%2819990315%2985:6<1353::AID-CNCR19>3.0.CO;2-1/full>
- A systematic review of the literature published in 2014 concluded that there was an association between exposure to glyphosate herbicides and non-Hodgkin's Lymphoma. <https://www.ncbi.nlm.nih.gov/pubmed/24762670>

Endocrine Disruption (Hormone Hacking)

- Glyphosate herbicide was a potent EDC in rats, causing disturbances in reproductive development after exposure during puberty. <https://www.ncbi.nlm.nih.gov/pubmed/20012598>
- This new Argentine study is the **first** to show endocrine-disrupting effects of a glyphosate-based herbicide on the uterus of newborn and pre-pubertal rats, supporting the possibility

that glyphosate-based herbicides are endocrine disruptors. <http://www.sciencedirect.com/science/article/pii/S0300483X16300932>

- An in vivo study of Roundup administered to rats in drinking water diluted to 50ng/L glyphosate equivalence – half of the level permitted in drinking water in the EU and 14,000 times lower than that permitted in drinking water in the USA – resulted in severe organ damage and a trend of increased incidence of mammary tumors in female animals over a 2-year period of exposure. The latter observation of tumors needs to be confirmed in an experiment with larger numbers of rats. <http://enveurope.springeropen.com/articles/10.1186/s12302-014-0014-5>

Kidney and Liver Damage at Low Doses

- A ground-breaking peer-reviewed study published in Environmental Health Journal in 2015 shows the levels of glyphosate-based herbicides which the general public are commonly exposed to in drinking water, altered the gene function of over 4000 genes in the livers and kidneys of rats. <http://ehjournal.biomedcentral.com/articles/10.1186/s12940-015-0056-1>

Binding of Vital Nutrients

- Glyphosate binds (chelates) vital nutrients such as iron, manganese, zinc, and boron in the soil, preventing plants from taking them up. <https://core.ac.uk/download/pdf/11741277.pdf?repositoryId=393>
- Genetically Modified (GM) soy plants treated with glyphosate have lower levels of essential nutrients and reduced growth, compared with GMO and non-GMO soy controls not treated with glyphosate. <http://link.springer.com/article/10.1007%2Fs11104-009-0081-3>

Antibiotic Resistance

- Research lead by a team from the University of Canterbury, New Zealand found that commonly used herbicides, including Roundup, can cause bacteria to become resistant to antibiotics. <http://mbio.asm.org/content/6/2/e00009-15>

For Reference: Allowed Levels of Glyphosate in Drinking Water

- Council of the European Union. Council directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption. Off J Eur Communities. 1998. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:1998:330:0032:0054:EN:PDF>
- US Environmental Protection Agency (EPA). Basic information about glyphosate in drinking water. 2014. <https://www.epa.gov/ground-water-and-drinking-water/table-regulated-drinking-water-contaminants>



Glyphosate Exposure Levels in Humans: Healthy and Chronically Diseased




Figure 3: Glyphosate in humans. A) Comparison of glyphosate excretion with urine of humans with conventional (N=99) and predominantly organic (N=41) feeding. B) Glyphosate in healthy (N=102) and chronically (N=199) diseased humans.

Source: Krüger M, 2014.



How to Avoid Glyphosate

After reading this report, which confirms glyphosate contamination in our food supply, as well as findings of glyphosate in drinking water, rain and the air, the first question you might ask yourself, is how can I avoid unwanted glyphosate residues in my, or my family's, daily meals?

The results presented in this report are the first round of independent testing of common American food products. This report uses the regulatory recognized "gold standard" testing methods of an FDA registered laboratory. While we are alarmed at the results found in a wide range of popularly consumed foods, we simply cannot say with certainty what levels of glyphosate residue are actually in the food supply.

These unprecedented pesticide residue tests confirm that glyphosate contamination is widespread. Not only do high levels of glyphosate residues show up in obvious GMO products, but they are also present in foods that are not genetically engineered. Glyphosate residues are now found in food crops, such as wheat, oats and barley, where glyphosate is used as a pre-harvest drying agent.

High Glyphosate Levels as a Result of Pre-Harvest Spraying of Roundup

In this initial round of testing, the two highest glyphosate residue levels were found on products that intentionally do not contain GMO ingredients and proudly boast their Non-GMO status on the packaging.

In the case of Cheerios, General Mills removed GMOs from their iconic cereal in 2014 and now market

the popular cereal as "Not made with genetically engineered ingredients" and also "gluten free." Unfortunately, the practice of pre-harvest spraying of Roundup on Cheerios number one ingredient "whole grain oats," has resulted in the highest levels of glyphosate contamination, which poses health concerns for young children consuming this food on a daily basis.

The next highest level was found in Stacy's Simply Naked Pita Chips, which contain no GMO ingredients and are actually certified by a third party, the Non-GMO Project, which test for GMO contamination levels, but not pesticide residues.

This report reveals that glyphosate use is widespread and moves freely in the environment. Even the two organic products that were tested as controls found glyphosate contamination. Obviously more testing is needed.

An Organic Diet for 1 Week Reduces Pesticide Exposures by 90%

Since GMOs and Roundup are both prohibited from organic production, the simplest way to avoid glyphosate contamination in food and to reduce exposures to synthetic pesticides is to eat organic food.

A 2014 study published in the Journal of Environmental Research confirmed that families eating an organic diet for as little as a week removed more than 90% of the pesticides from their system, as detected through urine tests.¹⁴ According to this independent study, "the average person is exposed to 10 to 13 pesticide residues each day from food, beverages and drinking water."

Call to Action:

Based on these scientific findings, Food Democracy Now! is calling for:

1. A federal investigation into the likely harmful effects of glyphosate on human health and the environment as a result of these disturbingly high levels of glyphosate residues found in popular American food products.
2. The EPA to refuse to reapprove glyphosate until the most current scientific evidence can be reviewed in an open and transparent process.
3. A permanent ban on the use of glyphosate as a pre-harvest drying agent for crops such as dry beans, sunflowers, wheat, oats and barley.
4. The immediate release of all restricted, allegedly "trade secret" data from all previous industry studies on glyphosate and glyphosate-based herbicides by the relevant federal agencies, including the Environmental Protection Agency (EPA), the Food and Drug Administration (FDA) and the U.S. Department of Agriculture (USDA).

Demand Immediate Release of Industry Science Data; End Pre-Harvest Spraying on Wheat, Oats, Barley and other Food Crops!

Today FOOD DEMOCRACY NOW! is calling for an immediate release by the EPA, FDA and USDA of all the data from corporate controlled scientific studies. Monsanto and other pesticide manufacturers are legally allowed to keep this information from the public and scientific community at large, based on the notion of "trade secrets," which since glyphosate's original patent expired is no longer a valid justification.

This lack of scientific integrity is alarming. Hiding fundamental research data from the public and the scientific community is a threat to public health. It undermines trust in the safety of our food supply, our federal government and the regulatory agencies that are supposed to be looking out for the wellbeing of the American public.

Urgent Need for Fundamental Reforms of Scientific Review Process

U.S. and European regulators and the chemical companies that supply them with corporate-sponsored research may try to dismiss these findings. However, a growing body of new independent, peer-reviewed scientific research continues to show disturbing evidence of harm from Roundup and glyphosate, at what were previously considered safe or extremely low doses. The evidence shows that Roundup and glyphosate are

far more toxic than was generally believed during the original scientific reviews of glyphosate's safety in the 1970s and 1980s.

Since that time, new scientific understandings of the real harm that chemicals can cause at low levels, such as the toxic effects of endocrine disruption, has emerged and must be considered in any new review of Roundup and glyphosate re-authorizations.



For comments on this report please contact:

David Murphy,
Executive Director, Food Democracy Now!
www.fooddemocracynow.org, dave@fooddemocracynow.org

Henry Rowlands,
Director, The Detox Project
www.detoxproject.org, henry@detoxproject.org

Food Democracy Now!: www.fooddemocracynow.org

Food Democracy Now! is a grassroots movement of more than 650,000 farmers and citizens dedicated to building a sustainable food system that protects our natural environment, sustains farmers and nourishes families.

We know we can build a food system that gives our communities equal access to healthy food, and respects the dignity of the farmers who produce it. We believe in recreating regional food systems, supporting the growth of humane, natural and organic farms, and protecting the environment. We value our children's health, worker's rights, conservation, and animal welfare over corporate profits. We believe that working together, we can make this vision a reality in our lifetimes.

The Detox Project: www.detoxproject.org

The Detox Project is a research platform that brings awareness to the public by testing for man-made chemicals in our bodies and in our food at a very personal level.

We believe you have the right to know what man-made chemicals are in your body and in your food!



About the Authors

David Murphy

Dave is the founder and executive director of Food Democracy Now!, a grassroots movement of more than 650,000 American farmers and citizens dedicated to reforming policies relating to food, agriculture and the environment.

In 2006, Murphy moved back to Iowa to help stop a factory farm from being built near his sister's farm. After seeing the loss of basic democratic rights of rural Iowans, Murphy decided to stay in Iowa to fight for Iowa's farmers and rural residents and expose the laws of industrial agriculture to help create a more sustainable future for all Americans.

In 2012, Murphy served as the co-chair of California's Prop 37, a ballot initiative to label genetically engineered foods. Following a narrow loss, Murphy helped write and pass the first two GMO labeling bills in Connecticut and Maine in 2013. His writing has appeared in *The Nation*, *The Hill*, *The Huffington Post* and *The New York Times*.

Food Democracy Now! Glyphosate: Unsafe on Any Plate

Henry Rowlands

Henry was brought up on a family run organic sheep farm in the Pembrokeshire National Park in Wales. His connection to both farming and the protection of the Welsh countryside led to a deep interest in issues related to sustainable agriculture from a young age.

Following work as a Journalist in many countries across Europe, Henry moved on to set up one of the World's most successful Sustainable Agriculture online news sources "Sustainable Pulse" which focuses on GMOs and pesticides. Sustainable Pulse now has a regular readership of over 100,000 people per month from over 125 countries.

Sustainable Pulse is also involved in a number of reference projects, all of which have the aim of educating the public on the problems surrounding the overuse of pesticides. These include The Detox Project, which has set up a unique pesticide testing platform across America.

References

- 1 "Acreage for Genetically Modified Crops Declined in 2015," The New York Times, April 13, 2016 <http://www.nytimes.com/2016/04/13/business/acreage-for-genetically-modified-crops-declined-in-2015.html>.
- 2 Mesnage R, Arno M, Costanzo M, Malatesta M, Seralini GE, Antoniou MN. Transcriptome profile analysis reflects rat liver and kidney damage following chronic ultra-low dose Roundup exposure. *Environ Health*. 2015;14:70. <http://ehjournal.biomedcentral.com/articles/10.1186/s12940-015-0056-1>.
- 3 Uren Webster TM, Santos EM. Global transcriptomic profiling demonstrates induction of oxidative stress and of compensatory cellular stress responses in brown trout exposed to glyphosate and Roundup. *BMC Genomics* 2015 Jan 31;16:32. PMID: 25636363 <http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-015-1254-5>.
- 4 Larsen K, Najle R, Lifschitz A, Virkel G. Effects of sub-lethal exposure of rats to the herbicide glyphosate in drinking water: glutathione transferase enzyme activities, levels of reduced glutathione and lipid peroxidation in liver, kidneys and small intestine. *Environ Toxicol Pharmacol*. 2012;34:811-8. doi: 10.1016/j.etap.2012.09.005. <https://www.ncbi.nlm.nih.gov/pubmed/23044091>.
- 5 Uren Webster TM, Santos EM. Global transcriptomic profiling demonstrates induction of oxidative stress and of compensatory cellular stress responses in brown trout exposed to glyphosate and Roundup. *BMC Genomics* 2015 Jan 31;16:32. PMID: 25636363 <http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-015-1254-5>.
- 6 Antoniou M et al. Teratogenic effects of glyphosate-based herbicides: Divergence of regulatory decisions from scientific evidence. *Journal of Environmental and Analytical Toxicology* 2012, 5:4. <http://www.omicsonline.org/teratogenic-effects-of-glyphosate-based-herbicides-divergence-of-regulatory-decisions-from-scientific-evidence-2161-0525.S4-006.pdf>
- 7 United States Patent 3,160,632 (1964) Stauffer Chemical: <http://1.usa.gov/1BULTJj>
- 8 United States Patent 3,799,758 (1974) Franz, Assignee Monsanto: <http://1.usa.gov/1BZlu02>
- 9 "How Much Money Does Monsanto Make From Roundup?," The Motley Fool, May 26, 2016. <http://www.fool.com/investing/2016/05/26/how-much-money-does-monsanto-make-from-roundup.aspx>.
- 10 Cakmak I, Yazici A, Tutus Y, Ozturk L. Glyphosate reduced seed and leaf concentrations of calcium, manganese, magnesium, and iron in non-glyphosate resistant soybean. *Eur J Agron*. 2009;31:114-119.
- 11 Neumann G, Kohls S, Landsberg E, Stock-Oliveira Souza K, Yamada T, Romheld V. Relevance of glyphosate transfer to non-target plants via the rhizosphere. *J Plant Dis Prot*. 2006;20:963-969.
- 12 Huber DM. What about glyphosate-induced manganese deficiency? *Fluid J*. 2007;20-22.
- 13 Bott S, Tesfamariam T, Candan H, Cakmak I, Romheld V, Neumann G. Glyphosate-induced impairment of plant growth and micronutrient status in glyphosate-resistant soybean (*Glycine max* L.). *Plant Soil*. 2008;312(1-2):185-194. doi:10.1007/s11104-008-9760-8.
- 14 Zobiolo LH, de Oliveira RS, Visentainer JV, Kremer RJ, Bellaloui N, Yamada T. Glyphosate affects seed composition in glyphosate-resistant soybean. *J Agric Food Chem*. 2010;58:4517-22. doi:10.1021/jf904342t.
- 15 Zobiolo LHS, de Oliveira RS, Huber DM, et al. Glyphosate reduces shoot concentrations of mineral nutrients in glyphosate-resistant soybeans. *Plant Soil*. 2010;328:57-69.
- 16 Kremer, R. J.; Means, N. E. Glyphosate and glyphosate resistant crop interactions with rhizosphere microorganisms. *Eur. J. Agron*. 2009, 31, 153-161.
- 17 Krüger M, Schrödl W, Neuhaus J, Shehata AA. Field investigations of glyphosate in urine of Danish dairy cows. *J Env Anal Toxicol*. 2013;3(5). doi: <http://dx.doi.org/10.4172/2161-0525.1000186>.
- 18 United States Patent 7,771,736 (2010) Abraham, Assignee Monsanto: <http://1.usa.gov/1EMmWz>
- 19 Benbrook, C. Trends in the use of glyphosate herbicide in the U.S. and globally. *Environmental Sciences Europe*. 2015;28(3). <http://enveurope.springeropen.com/articles/10.1186/s12302-016-0070-0>.
- 20 "Glyphosate Now the Most-Used Agricultural Chemical Ever," February 2, 2016, Newsweek <http://www.newsweek.com/glyphosate-now-most-used-agricultural-chemical-ever-422419>.
- 21 Glyphosate Map of America, Detox Project, Estimated Agricultural Use 1992 thru 2012. Source: USGS, Pesticide National Synthesis Project., <http://detoxproject.org/glyphosate-map-of-america/>.
- 22 National Water-Quality Assessment (NAWQA) Program, Pesticide National Synthesis Project, Estimated Agricultural Use for Glyphosate 1992 to 2013. https://water.usgs.gov/nawqa/pnsp/usage/maps/show_map.php?year=2013&map=GLYPHOSATE&hilo=L.
- 23 USDA ERS (2015). Adoption of genetically engineered crops in the U.S. USDA Economic Research Service. <http://www.ers.usda.gov/data-products/adoption-of-genetically-engineered-crops-in-the-us/recent-trends-in-ge-adoption.aspx>.
- 24 "U.S. researchers find Roundup chemical in water, air," August 31, 2011, Reuters <http://www.reuters.com/article/us-glyphosate-pollution-idUSTRE77U61720110831>.
- 25 Chang FC, Simcik MF, Capel PD. 2011. Occurrence and fate of the herbicide glyphosate and its degradate aminomethylphosphonic acid in the atmosphere. *Environ Toxicol Chem* 30:548-555. <http://dx.doi.org/10.1002/etc.431>.
- 26 Coupe RH, Capel PD: Trends in pesticide use on soybean, corn and cotton since the introduction of major genetically modified crops in the United States. *Pest Manag Sci* 2015.
- 27 "EPA tosses aside safety data, says Dow pesticide for GMOs won't harm people," December 8, 2015, Chicago Tribune. <http://www.chicagotribune.com/news/watchdog/ct-gmo-crops-pesticide-resistance-met-20151203-story.html>.
- 28 "Why is Glyphosate Sprayed on Crops Right Before Harvest?," March 5, 2016, EcoWatch. <http://www.ecowatch.com/why-is-glyphosate-sprayed-on-crops-right-before-harvest-1882187755.html>.
- 29 Hird, S.J. et al. Liquid chromatography-mass spectrometry for the determination of chemical contaminants in food, *TrAC Trends Anal. Chem*. 59 (2013) 59-72. <http://www.sciencedirect.com/science/article/pii/S0165993614000971>.
- 30 "Glyphosate Testing Revolution – The New Science, Sustainable Pulse," April 20, 2016. http://sustainablepulse.com/2016/04/20/a-guide-to-the-glyphosate-testing-revolution/#.V_mePWZSk1.
- 31 ResearchGate, Best method to detect pesticide (exposure) in blood samples, July 15, 2014. https://www.researchgate.net/post/Can_anyone_suggest_which_is_the_best_method_to_detect_pesticide_exposure_in_blood_samples.
- 32 "FDA to Start Testing for Glyphosate in Food," February 17, 2016, Time. <http://time.com/4227500/fda-glyphosate-testing/>.
- 33 "Roundup Chemical Glyphosate Found in 93% of Americans," About Lawsuits, June 1, 2016 <http://www.aboutlawsuits.com/roundup-chemical-glyphosate-testing-102183/>.
- 34 "Greens warn: German breast milk unsafe," June 26, 2016, The Local de. <http://www.thelocal.de/20160626/concerns-over-safety-of-german-breast-milk>.
- 35 "Cancer-linked pesticide found in popular German beer," February 26, 2016, RT. <https://www.rt.com/news/333679-cancer-pesticide-beer-germany/>.
- 36 "85% of Tampons Contain Monsanto's 'Cancer Causing' Glyphosate," October 26, 2015. <http://www.ecowatch.com/85-of-tampons-contain-monsantos-cancer-causing-glyphosate-188212780.html>
- 37 "150 European Parliament Members to Test Urine for Glyphosate," EcoWatch, April 11, 2016. <http://www.ecowatch.com/150-european-parliament-members-to-test-urine-for-glyphosate-1891081633.html>.
- 38 (PRiF) UDECoPRiF: Monitoring program. <http://www.food.gov.uk/business-industry/farmingfood/pesticides>.
- 39 Agricultural Marketing Service. Pesticide data program annual summary, program year 2011. In: Appendix C Distribution of Residues in Soybean by Pesticide. Washington, D.C.: U.S. Department of Agriculture; 2013.
- 40 EPA, Drinking Water Contaminants – Standards and Regulations, Table of Regulated Drinking Water Contaminants, What are glyphosate's health effects?. <https://safewater.zendesk.com/hc/en-us/articles/21402278-3-What-are-glyphosate-s-health-effects->.
- 41 Mesnage R, Arno M, Costanzo M, Malatesta M, Seralini GE, Antoniou MN. Transcriptome profile analysis reflects rat liver and kidney damage following chronic ultra-low dose Roundup exposure. *Environ Health*. 2015;14:70. <http://ehjournal.biomedcentral.com/articles/10.1186/s12940-015-0056-1>.
- 42 Seralini GE, Clair E, Mesnage R, Gress S, Defarge N, Malatesta M, et al. Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. *Environ Sci Europe*. 2014;26:14. <http://enveurope.springeropen.com/articles/10.1186/s12302-014-0014-5>.

- 43 Uren Webster TM, Santos EM. Global transcriptomic profiling demonstrates induction of oxidative stress and of compensatory cellular stress responses in brown trout exposed to glyphosate and Roundup. *BMC Genomics* 2015 Jan 31;16:32. PMID: 25636363 <http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-015-1254-5>.
- 44 Larsen K, Najle R, Lifschitz A, Virkel G. Effects of sub-lethal exposure of rats to the herbicide glyphosate in drinking water: glutathione transferase enzyme activities, levels of reduced glutathione and lipid peroxidation in liver, kidneys and small intestine. *Environ Toxicol Pharmacol.* 2012;34:811-8. doi: 10.1016/j.etap.2012.09.005. <https://www.ncbi.nlm.nih.gov/pubmed/23044094>.
- 45 EPA. Drinking Water Contaminants – Standards and Regulations, Table of Regulated Drinking Water Contaminants. <https://www.epa.gov/ground-water-and-drinking-water/table-regulated-drinking-water-contaminants#Inorganic>.
- 46 "Great Glyphosate Rebellion Continues as Europe Refuses Temporary License Extension," Sustainable Pulse, June 6, 2016. http://sustainablepulse.com/2016/06/06/great-glyphosate-rebellion-continues-as-europe-refuses-temporary-license-extension/#_b6m2ZS161.
- 47 "U.S. lawsuits build against Monsanto over alleged Roundup cancer link," Reuters, October 15, 2015. <http://www.reuters.com/article/us-usa-monsanto-lawsuits-idUSKCN0S92H720151015>.
- 48 "What Killed Jack McCall? A California Farmer Dies and a Case Against Monsanto Takes Root," *Huffington Post*, May 6, 2016 http://www.huffingtonpost.com/carey-gilliam/what-killed-jack-mccall-a_b_9852276.html.
- 49 Monsanto Roundup Lawsuit, Baum, Hedlund, Aristei and Goldman. <https://www.baumhedlundlaw.com/toxic-tort-law/monsanto-roundup-lawsuit/>.
- 50 "Health costs of hormone disrupting chemicals over € 150 bn a year in Europe, says study," *The UK Guardian*, March 6, 2015. <https://www.theguardian.com/environment/2015/mar/06/health-costs-hormone-disrupting-chemicals-150-bn-a-year-europe-says-study>.
- 51 Cakmak, I.; Yazici, A.; Tutus, Y.; Ozturk, L. Glyphosate reduced seed and leaf concentrations of calcium, manganese, magnesium, and iron in non-glyphosate resistant soybean. *Eur. J. Agron.* 2009, 31, 114–119. <http://research.sabanciuniv.edu/13147/>.
- 52 Kurenbach B, Marjoshi D, Amabile-Cuevas CF, Ferguson GC, Godsoe W, Gibson P, Heinemann JA et al. Sublethal exposure to commercial formulations of the herbicides dicamba, 2,4-dichlorophenoxyacetic acid, and glyphosate cause changes in antibiotic susceptibility in *Escherichia coli* and *Salmonella enterica* serovar typhimurium. *mBio.* 2015;6:2. <http://mbio.asm.org/content/6/2/e00009-15>.
- 53 Glyphosate: not JUST a carcinogen, *Wheat Belly Blog*, Dr. William Davis, October 12, 2015 <http://www.wheatbellyblog.com/2015/10/glyphosate-not-just-a-carcinogen/>.
- 54 Krüger M, Shehata AA, Schrödl W, Rodloff A. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe* 2013;20:74–78. <https://www.ncbi.nlm.nih.gov/pubmed/23396248>.
- 55 Shehata AA, Schrödl W, Aldin AA, Hafez HM, Krüger M. The effect of glyphosate on potential pathogens and beneficial members of poultry microbiota in vitro. *Curr Microbiol.* 2013;66(4):350–8. <http://link.springer.com/article/10.1007/s200284-012-0277-2>.
- 56 Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs DR, Lee DH, et al. Regulatory decisions on endocrine disrupting chemicals should be based on the principles of endocrinology. *Reprod Toxicol.* 2013;38C:1–15. <https://www.ncbi.nlm.nih.gov/pubmed/22419778>.
- 57 Thongprakaisang S, Thiantanawat A, Rangkadilok N, Suriyo T, Sata-yavivad J. Glyphosate induces human breast cancer cells growth via estrogen receptors. *Food Chem Toxicol.* 2013;59C:129–36. <http://euro-pepmc.org/abstract/med/23756170>.
- 58 Seralini GE, Clair E, Mesnage R, Gress S, Defarge N, Malatesta M, et al. Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. *Environ Sci Europe.* 2014;26:14. <https://enveurope.springeropen.com/articles/10.1186/s12302-014-0014-5>.
- 59 Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs DR, Lee DH, et al. Regulatory decisions on endocrine disrupting chemicals should be based on the principles of endocrinology. *Reprod Toxicol.* 2013;38C:1–15. <https://www.ncbi.nlm.nih.gov/pubmed/22419778>.
- 60 Mesnage R, Bernay B, Seralini GE. Ethoxylated adjuvants of glyphosate-based herbicides are active principles of human cell toxicity. *Toxicology.* 2013;313(2–3):122–8. <http://europepmc.org/abstract/MED/23000283>.
- 61 Antoniou M, Habib MEM, Howard CV, et al. Teratogenic effects of glyphosate-based herbicides: Divergence of regulatory decisions from scientific evidence. *J Env Anal Toxicol.* 2012;34:006. doi:10.4172/2161-0525.S4-006.
- 62 Romano RM, Romano MA, Bernardi MM, Furtado PV, Oliveira CA. Prepubertal exposure to commercial formulation of the herbicide Glyphosate alters testosterone levels and testicular morphology. *Arch Toxicol.* 2010;84:309-317.
- 63 Benedetti AL, Vituri C de L, Trentin AG, Domingues MA, Alvarez-Silva M. The effects of sub-chronic exposure of Wistar rats to the herbicide Glyphosate-Biocarb. *Toxicol Lett.* 2004;153:227-232. doi:10.1016/j.toxlet.2004.04.008.
- 64 Lee H-L, Kan C-D, Tsai C-L, Liou M-J, Guo H-R. Comparative effects of the formulation of glyphosate-surfactant herbicides on hemodynamics in swine. *Clin Toxicol Phila Pa.* 2009;47(7):651-658. doi:10.1080/15563650903158862. <https://www.ncbi.nlm.nih.gov/pubmed/22787363>.
- 65 Adam A, Marzuki A, Abdul Rahman H, Abdul Aziz M. The oral and intratracheal toxicities of ROUNDUP and its components to rats. *Vet Hum Toxicol.* 1997;39(3):147-151. <https://www.ncbi.nlm.nih.gov/pubmed/9167243>.
- 66 "Roundup: Birth Defects Caused By World's Top-Selling Weed killer, Scientists Say," *Huffington Post*, June 24, 2011 http://www.huffingtonpost.com/2011/06/24/roundup-scientists-birth-defects_n_883578.html.
- 67 Antoniou M, Habib MEM, Howard CV, et al. Teratogenic effects of glyphosate-based herbicides: Divergence of regulatory decisions from scientific evidence. *J Env Anal Toxicol.* 2012;34:006. doi:10.4172/2161-0525.S4-006. <http://www.omicsonline.org/open-access/the-formation-of-doxorubicin-loaded-targeted-nanoparticles-using-nanoprecipitation-double-emulsion-and-single-emulsion-for-cancer-2157-7439-1000379.php?aid=74534>.
- 68 Rapporteur member state, Germany. 1998. Monograph on Glyphosate. Released by the German Federal Agency for Consumer Protection and Food Safety, BVL. Volume 3-1, Glyphosat_05.pdf <https://www.scribd.com/document/57155616/VOLUME3-1-GLYPHOSAT-05>.
- 69 Larsen K, Najle R, Lifschitz A, Virkel G. Effects of sub-lethal exposure of rats to the herbicide glyphosate in drinking water: glutathione transferase enzyme activities, levels of reduced glutathione and lipid peroxidation in liver, kidneys and small intestine. *Environ Toxicol Pharmacol.* 2012;34(3):811-818. doi:10.1016/j.etap.2012.09.005. <https://www.ncbi.nlm.nih.gov/pubmed/23044094>.
- 70 Thongprakaisang S, Thiantanawat A, Rangkadilok N, Suriyo T, Sata-yavivad J. Glyphosate induces human breast cancer cells growth via estrogen receptors. *Food Chem Toxicol.* 2013;59:129–136. doi:10.1016/j.fct.2013.05.057. <https://www.ncbi.nlm.nih.gov/pubmed/23756170>.
- 71 Attorney General of the State of New York 1996. In the Matter of Monsanto Company, Respondent. Assurance of Discontinuance Pursuant to Executive Law § 63(15). New York: Attorney General of the State of New York, Consumer Frauds and Protection Bureau, Environmental Protection Bureau.
- 72 EPA 40 CFR Part 180 [EPA-HQ-OPP-2012-0132; FRL-9384-3] Glyphosate; Pesticide Tolerances, Federal Register/ Vol. 78, No. 84 / Wednesday, May 1, 2013 / Rules and Regulations <https://www.gpo.gov/fdsys/pkg/FR-2013-05-01/pdf/2013-10316.pdf>.
- 73 Pesticide Tolerances: Glyphosate, Rule document issued by Environmental Protection Agency, Regulations dot gov, Comment Period Closed July 12 2013. <https://www.regulations.gov/document?D=EPA-HQ-OPP-2012-0132-0009>.
- 74 Benbrook C. Trends in the use of glyphosate herbicide in the U.S. and globally. *Environmental Sciences Europe.* 2015;28(3). <http://dx.doi.org/10.1186/s12302-015-0070-0>.
- 75 "EPA raises levels of glyphosate residue allowed in food," July 5, 2013, *The Washington Times*. <https://web.archive.org/web/20130709080009/http://communities.washingtontimes.com/neighborhood/world-our-backyard/2013/jul/5/epa-raises-levels-glyphosate-residue-allowed-your->
- 76 Bohn T, Cuha M, Traavik T, Sanden M, Fagan J, Primicerio R (2014) Compositional differences in soybeans on the market: glyphosate accumulation in Roundup Ready GM soybeans. *Food Chem* 153:207–215. <http://www.sciencedirect.com/science/article/pii/S0308814613010201>
- 77 "How 'Extreme Levels' of Roundup in Food Became the Industry Norm," March 24, 2014, Thomas Bohn and Marek Cuhra, *Independent Science*. <http://www.independentsciencenews.org/news/how-extreme-levels-of-roundup-in-food-became-the-industry-norm/>

- 78 "UCSF Presentation Reveals Glyphosate Contamination in People across America," May 25, 2016, The Detox Project. <http://detoxproject.org/13212/>
- 79 "Glyphosate Found in Urine of 93 Percent of Americans Tested," May 29, 2016, EcoWatch. <http://www.ecowatch.com/glyphosate-found-in-urine-of-93-percent-of-americans-tested-189146755.html>
- 80 Krüger M, Schledorn P, Schrödl W, Hoppe HW, Lutz W, et al. (2014) Detection of Glyphosate Residues in Animals and Humans. *J Environ Anal Toxicol* 4: 210.
- 81 WHO (1987). "Principles for the safety assessment of food additives and contaminants in food." *Environmental Health Criteria* 70 <http://www.inchem.org/documents/ehc/ehc/ehc70.htm#SectionNumber:5.5>
- 82 Monograph on Glyphosate. Annex B-5.10.2: Toxicology and Metabolism (1998) Released by German government agency BVL. www.scribd.com/doc/57155616/VOLUME3-1-GLYPHOSAT-05
- 83 For EPA's setting of the glyphosate ADI at 0.1mg/kg/day in the early 1980s (vs. 175 today), see EPA (1983). Glyphosate (Roundup) on wheat. March 3, 1983.
- 84 "Monsanto Knew of Glyphosate Cancer Link 35 Years Ago," Sustainable Pulse, April 9, 2015 <http://sustainablepulse.com/2015/04/09/monsanto-knew-of-glyphosate-cancer-link-35-years-ago/>
- 85 EPA SECOND Peer Review of Glyphosate, The Health Effects Division (HED) Carcinogenicity Peer Review Committee (CPRC) June 26, 1991.
- 86 Rapporteur member state, Germany. 1998. Monograph on Glyphosate. Released by the German Federal Agency for Consumer Protection and Food Safety, BVL. Volume 3-1_Glyphosat_05.pdf <https://www.scribd.com/document/57155616/VOLUME3-1-GLYPHOSAT-05>
- 87 Rapporteur member state, Germany. 1998. Monograph on Glyphosate. Released by the German Federal Agency for Consumer Protection and Food Safety, BVL. Volume 3-1_Glyphosat_05.pdf <https://www.scribd.com/document/57155616/VOLUME3-1-GLYPHOSAT-05>
- 88 For EPA's setting of the glyphosate ADI at 0.1mg/kg/day in the early 1980s (vs. 175 today), see EPA (1983). Glyphosate (Roundup) on wheat. March 3, 1983.
- 89 Rapporteur member state, Germany. 1998. Monograph on Glyphosate. Released by the German Federal Agency for Consumer Protection and Food Safety, BVL. Volume 3-1_Glyphosat_05.pdf <https://www.scribd.com/document/57155616/VOLUME3-1-GLYPHOSAT-05>
- 90 Mesnage et al. Transcriptome profile analysis reflects rat liver and kidney damage following chronic ultra-low dose Roundup exposure. *Environmental Health* 2015;14:70. <https://ehjournal.biomedcentral.com/articles/10.1186/s12940-015-0056-1>
- 91 Antoniou M et al. Teratogenic effects of glyphosate-based herbicides: Divergence of regulatory decisions from scientific evidence. *Journal of Environmental and Analytical Toxicology* 2012, S:4. <http://www.omicsonline.org/teratogenic-effects-of-glyphosate-based-herbicides-divergence-of-regulatory-decisions-from-scientific-evidence-2161-0525-S4-006.pdf>
- 92 "Gene expression analysis confirms Roundup causes liver and kidney damage at very low doses," GMWatch, August 25, 2015. <http://www.gmwatch.org/news/latest-news/16375-gene-expression-analysis-confirms-roundup-causes-liver-and-kidney-damage-at-very-low-doses>.
- 93 Monsanto Europe, December 1995.
- 94 Richard S, Moslemi S, Sipahutar H, Benachour N, Seralini G-E (2005) Differential effects of glyphosate and roundup on human placental cells and aromatase. *Environ Health Perspect* 113:716-720. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1257596/>.
- 95 Mesnage R, Defarge N, Spiroux de Vendômois J, Seralini GE (2014) Major pesticides are more toxic to human cells than their declared active principles. *Biomed Res Int* 2014:179691 <https://www.hindawi.com/journals/bmri/2014/179691/citations/>.
- 96 Johal, G.R. and Huber, D.M. 2009. Glyphosate effects on diseases of plants. *European J. Agron.* 31:144-152. <http://www.certifiedorganic.bc.ca/rcbtoa/services/huber-glyphosates-2009.pdf>.
- 97 "Scientist warns of dire consequences with widespread use of glyphosate," The Organic and Non-GMO Report, May 2010. http://www.non-gmoreport.com/articles/may10/consequenceso_widespread_glyphosate_use.php.
- 98 Defarge, N., Takács, E., Lozano, V. L., Mesnage, R., Spiroux de Vendômois, J., Seralini, G.-E., & Székács, A. (2016). Co-Formulants in Glyphosate-Based Herbicides Disrupt Aromatase Activity in Human Cells below Toxic Levels. *International Journal of Environmental Research and Public Health*, 13(3), 264. <http://doi.org/10.3390/ijerph13030264>.
- 99 "New Evidence About the Dangers of Monsanto's Roundup," The Intercept, May 17, 2016. <https://theintercept.com/2016/05/17/new-evidence-about-the-dangers-of-monsantos-roundup/>.
- 100 "Weed-Wacking Herbicide Proves Deadly to Human Cells," *Scientific American*, June 23, 2009. <http://www.scientificamerican.com/article/weed-whacking-herbicide-p/>.
- 101 Benachour, N., and G. E. Seralini. "Glyphosate Formulations Induce Apoptosis and Necrosis in Human Umbilical, Embryonic, and Placental Cells." [In eng]. *Chem Res Toxicol* 22, no. 1 (Jan 2009): 97-105. <https://www.ncbi.nlm.nih.gov/pubmed/19105591>
- 102 Krüger M, Schledorn P, Schrödl W, Hoppe HW, Lutz W, et al. (2014) Detection of Glyphosate Residues in Animals and Humans. *J Environ Anal Toxicol* 4: 210. <http://www.omicsonline.org/open-access/detection-of-glyphosate-residues-in-animals-and-humans-2161-0525-1000210.php?aid=23853>.
- 103 Monsanto, Background: Summary on Human Risk Assessment and Safety Evaluation on Glyphosate and Roundup Herbicide (Updated November 2014), Williams, Gary M, Kroes, Robert, Munro, Ian C. <http://www.monsanto.com/glyphosate/documents/summary-of-human-risk-assessment-and-safety-evaluation.pdf>.
- 104 USEPA. 1993. Reregistration Eligibility Decision (RED) Glyphosate. Office of Prevention, Pesticides and Toxic Substances. Washington DC.
- 105 Krüger M, Schledorn P, Schrödl W, Hoppe HW, Lutz W, et al. (2014) Detection of Glyphosate Residues in Animals and Humans. *J Environ Anal Toxicol* 4: 210.
- 106 Larsen K, Najle R, Lifschitz A, Mate ML, Lanusse C, Virkel GL. Effects of Sublethal Exposure to a Glyphosate-Based Herbicide Formulation on Metabolic Activities of Different Xenobiotic-Metabolizing Enzymes in Rats. *Int J Toxicol* 2014. <https://www.ncbi.nlm.nih.gov/pubmed/24985121>.
- 107 Franks AL, Slansky JE. Multiple associations between a broad spectrum of autoimmune diseases, chronic inflammatory diseases and cancer. *Anticancer Res.* 2012;32:119-36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3349285/eb>.
- 108 IARC Monograph Glyphosate is probably carcinogenic to humans (Group 2A) 2015, <http://monographs.iarc.fr/ENG/Monographs/vol112/mono112-09.pdf>.
- 109 Parks CG, Hoppin JA, DeRoos AJ, Costenbader KH, Alavanja MC, Sandler DP. 2016. Rheumatoid arthritis in Agricultural Health Study spouses: associations with pesticides and other farm exposures. *Environ Health Perspect*; doi:10.1289/EHP129 <http://ehp.niehs.nih.gov/ehp129/>.
- 110 "Glyphosate link with rheumatoid arthritis shown in new study," GMWatch, June 12, 2016 <http://www.gmwatch.org/news/latest-news/17022-glyphosate-link-with-rheumatoid-arthritis-shown-in-new-study>.
- 111 International Programme on Chemical Safety (IPCS), Pesticide Residues in Food 2004, Evaluations Part II Toxicological, Joint FAO/WHO Meeting on Pesticide Residues. http://apps.who.int/iris/bit-stream/10665/43624/1/9241665203_eng.pdf.
- 112 Attorney General of the State of New York 1996. In the Matter of Monsanto Company, Respondent. Assurance of Discontinuance Pursuant to Executive Law § 63(15). New York: Attorney General of the State of New York, Consumer Frauds and Protection Bureau, Environmental Protection Bureau.
- 113 "Monsanto Fined in France for 'False' Herbicide Ads." TerraDaily.com. January 27, 2007. <http://www.terradaily.com/2006/070126154451.ovopjxml.html>
- 114 Oates L, Cohen M, Braun L, Schembri A, Taskova R. Reduction in urinary organophosphate pesticide metabolites in adults after a week-long organic diet. *Environ Res.* 2014;132:105-111. <https://www.ncbi.nlm.nih.gov/pubmed/24769399>